

Demuškin Groups with Operators

by Kay Wingberg at Heidelberg

In this paper we consider pro- p Poincaré groups of dimension 2, also called “Demuškin groups”. These groups are defined as follows: a pro- p -group G is called a Demuškin group if its cohomology has the following properties:

$$\begin{aligned} \dim_{\mathbb{F}_p} H^1(G, \mathbb{Z}/p\mathbb{Z}) &< \infty, \\ \dim_{\mathbb{F}_p} H^2(G, \mathbb{Z}/p\mathbb{Z}) &= 1, \text{ and the cup-product} \\ H^1(G, \mathbb{Z}/p\mathbb{Z}) \times H^1(G, \mathbb{Z}/p\mathbb{Z}) &\xrightarrow{\cup} H^2(G, \mathbb{Z}/p\mathbb{Z}) \text{ is non-degenerate.} \end{aligned}$$

In the following we exclude the exceptional case that $G \cong \mathbb{Z}/2\mathbb{Z}$. The dualizing module I of G is isomorphic to $\mathbb{Q}_p/\mathbb{Z}_p$ as an abelian group and we have a canonical action of G on I .

Demuškin groups occur as Galois groups of the maximal p -extension of p -adic number fields (if these fields contain the group of p -th roots of unity) and as the p -completion of the fundamental group of a compact oriented Riemann surface. In the first case the action of G on I is non-trivial whereas in the second case G acts trivially on I . We will only consider Demuškin groups acting non-trivially on its dualizing module and we are interested in free pro- p -quotients of these groups. Possible ranks of such free quotients were first calculated in [7], [6] and [2].

In many cases of interest a finite group Δ of order prime to p acts on a Demuškin group. As an example consider the local field $k = \mathbb{Q}_p(\zeta_p)$, where p is an odd prime number. Then $G(k|\mathbb{Q}_p) \cong \mathbb{Z}/(p-1)\mathbb{Z}$ acts on the Demuškin group $D = G(k(p)|k)$, where $k(p)$ is the maximal p -extension of k . Of particular interest is the case where Δ is generated by an involution, e.g. $G(k|\mathbb{Q}_p(\zeta_p + \zeta_p^{-1})) \cong \mathbb{Z}/2\mathbb{Z}$ acts on D ; see [8] where Demuškin groups with involution were considered.

In the following, we are mainly interested in free pro- p -quotients of a Demuškin group G which are invariant under a given action of Δ on G . We show the existence of Δ -invariant free quotients F of G such that F^{ab} has a prescribed action of Δ .

If p is odd and $\Delta \cong \mathbb{Z}/2\mathbb{Z}$ acts on a p -Demuškin group G of rank $n + 2$, then there exists a Δ -invariant free quotient F of G such that $\text{rank}_{\mathbb{Z}_p}(F^{ab})^+ = 1$ and $\text{rank}_{\mathbb{Z}_p}(F^{ab})^- = n/2$. This situation occurs for number fields as the following example shows:

Let p be an odd regular prime number and consider the CM-field $k = \mathbb{Q}(\zeta_p)$ with maximal totally real subfield $k^+ = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$. Then the Galois group $G(k|k^+) \cong \mathbb{Z}/2\mathbb{Z}$ acts on the Galois group $G(k_{S_p}(p)|k)$ of the maximal p -extension $k_{S_p}(p)$ of k which is unramified outside p . Let $k_{\mathfrak{p}}$ be the completion of k with respect to the unique prime $\mathfrak{p}$ of k above p and let $k_{\mathfrak{p}}(p)$ its maximal p -extension. Then we have a surjection

$$G(k_{\mathfrak{p}}(p)|k_{\mathfrak{p}}) \quad G(k_{S_p}(p)|k)$$

of the Demuškin group $G(k_{\mathfrak{p}}(p)|k_{\mathfrak{p}})$ of rank $p + 1$ onto the free pro- p -group $G(k_{S_p}(p)|k)$ of rank $(p + 1)/2$ and

$$\text{rank}_{\mathbb{Z}_p}(G(k_{S_p}(p)|k)^{ab})^+ = 1, \quad \text{rank}_{\mathbb{Z}_p}(G(k_{S_p}(p)|k)^{ab})^- = (p - 1)/2.$$

It would be of interest under which conditions there exist large free quotients of $G(k_{S_p}(p)|k)$ for an arbitrary CM-field k . If we assume that no prime $\mathfrak{p}$ above p splits in the extension $k|k^+$, such a quotient should be defined by free quotients of the local groups $G(k_{\mathfrak{p}}(p)|k_{\mathfrak{p}})$, $\mathfrak{p}|p$, with an acting of $G(k_{\mathfrak{p}}|k_p^+) \cong G(k|k^+)$ as above.

1 Operators

Let p be a prime number and let

$$1 \quad G \quad \overset{s}{\mathcal{G}} \quad \Delta \quad 1,$$

be a split exact sequence of profinite groups where G is a pro- p -group and Δ is a finite group of order prime to p . Thus $\mathcal{G}$ is the semi-direct product of Δ by G and G is a pro- p - Δ operator group where the action of Δ on G is defined via the splitting s . Conversely, given a pro- p - Δ operator group G , we get a semi-direct product $\mathcal{G} = G \rtimes \Delta$ where the action of Δ on G is the given one.

Let $\mathcal{G}(p)$ be the maximal pro- p -quotient of $\mathcal{G}$ and let G_{Δ} be the maximal quotient of G with trivial Δ -action. Observe that G_{Δ} is well-defined.

Proposition 1.1 *With the notation and assumptions as above there is a canonical isomorphism*

$$G_{\Delta} \xrightarrow{\sim} \mathcal{G}(p).$$

Furthermore, if $H^2(G, \mathbb{Z}/p\mathbb{Z})^{\Delta} = 0$, then G_{Δ} is a free pro- p -group.

Proof: Consider the exact commutative diagram

$$\begin{array}{ccccccc}
& & & s & & & \\
1 & & N \cap G & \longrightarrow & N & \xrightarrow{s} & \Delta & \longrightarrow & 1 \\
& & & & & & & & \\
1 & & G & \longrightarrow & \mathcal{G} & \xrightarrow{s} & \Delta & \longrightarrow & 1 \\
& & & & & & & & \\
& & \tilde{G} & \longrightarrow & \mathcal{G}(p), & & & &
\end{array}$$

where N is the kernel of the canonical surjection $\mathcal{G} \rightarrow \mathcal{G}(p)$ and $\tilde{G}$ denotes the quotient $G/N \cap G$. Since Δ acts on $N \cap G$ via s , we obtain an induced action on $\tilde{G}$. This action is trivial because

$$g^{s(\sigma)-1} = [s(\sigma), g] \in N \cap G \quad \text{for } g \in G \text{ and } \sigma \in \Delta,$$

and so we get a surjection

$$\varphi : G_\Delta \twoheadrightarrow \tilde{G}.$$

Consider the exact commutative diagram

$$\begin{array}{ccccccc}
0 & & H^1(\tilde{G}) & \longrightarrow & H^1(G_\Delta) & \longrightarrow & H^1(\ker \varphi)^{G_\Delta} & \longrightarrow & H^2(\tilde{G}) & \longrightarrow & H^2(G_\Delta) \\
& & & & \text{\textit{inf}} & & & & & & \text{\textit{inf}} \\
& & \text{\textit{res}} & & H^1(G)^\Delta & & \text{\textit{res}} & & H^2(G)^\Delta & & \text{\textit{res}} \\
& & & & \text{\textit{res}} & & & & & & \\
H^1(\mathcal{G}(p)) & \xrightarrow{\text{\textit{inf}}_1} & H^1(\mathcal{G}) & & & & H^2(\mathcal{G}(p)) & \xrightarrow{\text{\textit{inf}}_2} & H^2(\mathcal{G})
\end{array}$$

where $H^i(-) = H^i(-, \mathbb{Z}/p\mathbb{Z})$, and the bijectivity of $\text{\textit{inf}}_1$ and the injectivity of $\text{\textit{inf}}_2$ follows from $\text{Hom}(N, \mathbb{Z}/p\mathbb{Z}) = 0$. We see that $H^1(\ker \varphi)^{G_\Delta} = 0$, and so $\ker \varphi = 1$, i.e. $G_\Delta \cong \tilde{G} \cong \mathcal{G}(p)$.

Furthermore, it follows that

$$H^2(G_\Delta) \xrightarrow{\text{\textit{inf}}} H^2(G)^\Delta$$

is injective. Therefore, if $H^2(G)^\Delta = 0$, then $H^2(G_\Delta) = 0$, and so G_Δ is a free pro- p -group. $\square$

In the following lemma let G be a finitely generated pro- p - Δ operator group, i.e. there is a homomorphism of profinite groups $\Delta \longrightarrow \text{Aut}(G)$, cf. [4] 5.1. Let

$$1 \longrightarrow R \longrightarrow F \longrightarrow G \longrightarrow 1$$

be a minimal presentation of the group G by a free pro- p -group F of rank $\dim_{\mathbb{F}_p} H^1(G, \mathbb{Z}/p\mathbb{Z})$ and a normal subgroup R .

Lemma 1.2 *With the notation as above there exists a continuous action of Δ on F extending the action on G , i.e. the surjection $F \twoheadrightarrow G$ is Δ -invariant.*

Proof: We may assume that the action of Δ on G is faithful, i.e. Δ injects into $\text{Aut}(G)$. Let $\sigma \in \Delta$. Since F is free, there exists a homomorphism $\tilde{\sigma} : F \rightarrow F$ such that the diagram

$$\begin{array}{ccc} & F & \\ \tilde{\sigma} & & \\ F & \xrightarrow{\sigma} & G \end{array}$$

commutes. The map $\tilde{\sigma}$ is necessarily an automorphism of F since it induces a bijection on $F/F^2 \cong G/G^2$.

Let $\tilde{\Delta}$ be the subgroup of $\text{Aut}(F)$ generated by all lifting $\tilde{\sigma}$ of the elements $\sigma \in \Delta$. Recall that the kernels of the homomorphisms $\text{Aut}(G) \rightarrow \text{Aut}(G/G^2)$ and $\text{Aut}(F) \twoheadrightarrow \text{Aut}(F/F^2)$ are pro- p -groups, cf. [4] 5.5, and that Δ has an order prime to p . Therefore, we can consider Δ as a subgroup of $\text{Aut}(G/G^2) \cong \text{Aut}(F/F^2)$, and there exists a subgroup Δ' of $\tilde{\Delta}$ such the diagram

$$\Delta' \subseteq \tilde{\Delta} \quad \text{Aut}(F)$$

$$\Delta \quad \text{Aut}(F/F^2)$$

commutes. Since every lifting $\tilde{\sigma}$ respects R , the same is true for the elements of Δ' , i.e. Δ' consists of liftings of elements of Δ . This proves the lemma. $\square$

Now we assume that

- Δ is a finite group of order prime to p and
- G is a p -Demuškin group of rank $n + 2, n \geq 0$, with dualizing module I and an action by Δ .

Let $\mathcal{G}$ be the semi-direct product of Δ by G , i.e. the sequence

$$1 \longrightarrow G \longrightarrow \mathcal{G} \longrightarrow \Delta \longrightarrow 1$$

is exact.

The dualizing module I of G is defined as

$$I = \varinjlim_m \varinjlim_U H^2(U, \mathbb{Z}/p^m\mathbb{Z})^*,$$

where U runs through the open normal subgroups of G and the second limit is taken over the maps cor^* dual to the corestriction; the first limit is taken with respect to the multiplication by p .

Let

$$\chi : G \longrightarrow \text{Aut}(I) \cong \mathbb{Z}_p^\times$$

be the character given by the action of G on I . We denote the canonical quotient $G/\ker(\chi)$ by Γ , i.e.

$$\chi_0 : \Gamma \longrightarrow \text{Aut}(I).$$

In the following we assume that

$$G \text{ acts non-trivially on } I$$

(thus $\Gamma \cong \mathbb{Z}_p$), and we define the (finite) invariant q of G by

$$q = \#(I^G).$$

Then we have a Δ -invariant isomorphism

$$H^2(G, \mathbb{Z}/q\mathbb{Z}) \cong \text{Hom}(I^G, \mathbb{Z}/q\mathbb{Z}) \quad (\cong \mathbb{Z}/q\mathbb{Z} \text{ as an abelian group})$$

and a Δ -invariant non-degenerate pairing

$$H^1(G, \mathbb{Z}/q\mathbb{Z}) \times H^1(G, \mathbb{Z}/q\mathbb{Z}) \xrightarrow{\cup} H^2(G, \mathbb{Z}/q\mathbb{Z}).$$

From the exact sequence $0 \rightarrow \mathbb{Z}/q\mathbb{Z} \xrightarrow{q} \mathbb{Z}/q^2\mathbb{Z} \rightarrow \mathbb{Z}/q\mathbb{Z} \rightarrow 0$, we get the Bockstein homomorphism

$$B : H^1(G, \mathbb{Z}/q\mathbb{Z}) \longrightarrow H^2(G, \mathbb{Z}/q\mathbb{Z})$$

which is surjective and Δ -invariant.

For a pro- p -group P we denote by P^i , $i \geq 1$, the descending q -central series, i.e.

$$P^1 = P \quad \text{and} \quad P^{i+1} = (P^i)^q [P^i, P] \quad \text{for } i \geq 1.$$

Let

$$1 \longrightarrow F \longrightarrow \mathcal{F} \longrightarrow \Delta \longrightarrow 1$$

be an exact sequence of profinite groups where F is a finitely generated pro- p -group. Obviously, G^i and F^i are normal open subgroups of $\mathcal{G}$ and $\mathcal{F}$ respectively.

Proposition 1.3 *With the notation as above let $q > 2$ and $m \geq 2$. Assume that there exists a surjection*

$$\varphi_{m+1} : \mathcal{G} \longrightarrow \mathcal{F}/F^{m+1}.$$

Then there exists a surjection

$$\varphi : \mathcal{G} \longrightarrow \mathcal{F}$$

inducing the surjection $\varphi_m : \mathcal{G} \xrightarrow{\varphi_{m+1}} \mathcal{F}/F^{m+1} \xrightarrow{\text{can}} \mathcal{F}/F^m$.

Proof: Assume that we have already found a surjection

$$\varphi_{i+1} : \mathcal{G} \twoheadrightarrow \mathcal{F}/F^{i+1}$$

for $i \geq m$ which induces φ_m , and let $\varphi_i : \mathcal{G} \xrightarrow{\varphi_{i+1}} \mathcal{F}/F^{i+1} \xrightarrow{\text{can}} \mathcal{F}/F^i$.

Let $\gamma, x_0, \dots, x_n$ be a minimal system of generators of G such that $x_k \in \ker(\chi)$ for $k \geq 0$ and $\chi(\gamma) = 1 - q$.

Claim: The group F^{i+1}/F^{i+2} is generated by elements of the form

$$w^q[w, \bar{\gamma}] \bmod F^{i+2}, \quad [w, \bar{x}_k] \bmod F^{i+2}, \quad k \geq 0, \quad w \in F^i,$$

where $\bar{\gamma}, \bar{x}_k \in F$ are lifts of $\varphi_2(\gamma), \varphi_2(x_k) \in F/F^2$.

This shown in [3] prop. 5(i) (observe, that we have a surjection $G/G^{i+1} \twoheadrightarrow F/F^{i+1}$, and so the group F/F^{i+1} is generated by the elements $\bar{\gamma}, \bar{x}_k \bmod F^{i+1}$).

Consider the diagram with exact line

$$\begin{array}{ccccccc}
 & & & \mathcal{G} & & & \\
 & & & \varphi_i & & & \\
 (*) & & & & & & \\
 & 1 & F^i/F^{i+2} & \mathcal{F}/F^{i+2} & \mathcal{F}/F^i & 1. &
 \end{array}$$

Since $i \geq m \geq 2$, the group F^i/F^{i+2} is abelian,

$$[F^i, F^i] \subseteq F^{2i} \subseteq F^{i+2},$$

and we consider F^i/F^{i+2} as a $\mathcal{G}$ -module via φ_i . Since the dualizing module I of G is isomorphic to $\mathbb{Q}_p/\mathbb{Z}_p$ as an abelian group, the canonical exact sequence

$$0 \longrightarrow F^{i+1}/F^{i+2} \longrightarrow F^i/F^{i+2} \longrightarrow F^i/F^{i+1} \longrightarrow 0$$

induces a Δ -invariant exact sequence

$$0 \longrightarrow \text{Hom}_G(F^i/F^{i+1}, I) \longrightarrow \text{Hom}_G(F^i/F^{i+2}, I) \longrightarrow \text{Hom}_G(F^{i+1}/F^{i+2}, I).$$

Let $f \in \text{Hom}_G(F^i/F^{i+2}, I)$. Then

$$\begin{aligned}
 f([w, \bar{x}_k] \bmod F^{i+2}) &= f(w \bmod F^{i+2})^{x_k-1} = 0 \quad \text{for } k \geq 0, \\
 f(w^q[w, \bar{\gamma}] \bmod F^{i+2}) &= f(w \bmod F^{i+2})q + f(w \bmod F^{i+2})^{\gamma-1} \\
 &= f(w \bmod F^{i+2})(q - q) = 0.
 \end{aligned}$$

Using the claim, we see that f vanishes on F^{i+1}/F^{i+2} , and so

$$\text{Hom}_G(F^i/F^{i+1}, I) \xrightarrow{\sim} \text{Hom}_G(F^i/F^{i+2}, I).$$

By duality, cf. [5] (3.7.6), (3.7.1), (3.4.6), we get

$$H^2(G, F^i/F^{i+2}) \xrightarrow{\sim} H^2(G, F^i/F^{i+1}),$$

and so

$$H^2(G, F^i/F^{i+2})^\Delta \xrightarrow{\sim} H^2(G, F^i/F^{i+1})^\Delta.$$

Since the order of Δ is prime to p , we obtain the isomorphism

$$H^2(\mathcal{G}, F^i/F^{i+2}) \xrightarrow{\sim} H^2(\mathcal{G}, F^i/F^{i+1}).$$

Now we prove that the embedding problem $(*)$ is solvable. For this we have to show that the 2-class

$$[\beta_i] \in H^2(\mathcal{F}/F^i, F^i/F^{i+2})$$

is mapped to zero under the inflation map $inf = \varphi_i^*$,

$$H^2(\mathcal{F}/F^i, F^i/F^{i+2}) \xrightarrow{inf} H^2(\mathcal{G}, F^i/F^{i+2}),$$

where β_i is the 2-cocycle corresponding to the group extension in $(*)$, see [5] (9.4.2). From the commutative exact diagram

$$\begin{array}{ccccccc} & & & \mathcal{G} & & & \\ & & & \varphi_i & & & \\ & & \varphi_{i+1} & & & & \\ 1 & F^i/F^{i+2} & \mathcal{F}/F^{i+2} & \mathcal{F}/F^i & 1 & \beta_i & \\ & can & & & & & \\ 1 & F^i/F^{i+1} & \mathcal{F}/F^{i+1} & \mathcal{F}/F^i & 1 & \alpha_i & \end{array}$$

we get a commutative diagram

$$\begin{array}{ccc} H^2(\mathcal{F}/F^i, F^i/F^{i+1}) & \xrightarrow{\varphi_i^*} & H^2(\mathcal{G}, F^i/F^{i+1}) \\ can_* & & can_* \\ H^2(\mathcal{F}/F^i, F^i/F^{i+2}) & \xrightarrow{\varphi_i^*} & H^2(\mathcal{G}, F^i/F^{i+2}). \end{array}$$

Since there exists the solution φ_{i+1} for the embedding problem α_i , we have $\varphi_i^*([\alpha_i]) = 0$, and so

$$can_* \circ \varphi_i^*([\beta_i]) = \varphi_i^* \circ can_*([\beta_i]) = \varphi_i^*([\alpha_i]) = 0.$$

From the injectivity of the map can_* on the right-hand side of the diagram above it follows that $\varphi_i^*([\beta_i]) = 0$, and so there exists a solution

$$\varphi_{i+2} : \mathcal{G} \longrightarrow \mathcal{F}/F^{i+2}$$

of the embedding problem corresponding to β_i . This homomorphism is necessarily surjective and induces φ_m , because φ_i has these properties, cf. [5] (3.9.2).

Using a compactness argument, we get in the limit a surjection $\varphi : \mathcal{G} \twoheadrightarrow \mathcal{F}$ inducing φ_m . This finishes the proof of the proposition. $\square$

In the following let p be an odd prime number and let $\Delta = \langle \sigma \rangle \cong \mathbb{Z}/2\mathbb{Z}$ be cyclic of order 2. We denote, as usual, the $(\pm)$ -eigenspaces of a $\mathbb{Z}_p[\Delta]$ -module M by $M^\pm$.

Proposition 1.4 *Let p be an odd prime number and let G be a p -Demuškin group of rank $n + 2$, $n \geq 0$, with dualizing module I and invariant $q = \#(I^G) < \infty$. Assume that $\Delta \cong \mathbb{Z}/2\mathbb{Z}$ acts on G . Then the following holds:*

- (i) *If $H^2(G, \mathbb{Z}/p\mathbb{Z}) = H^2(G, \mathbb{Z}/p\mathbb{Z})^-$, then G_Δ is a free pro- p -group of rank $n/2 + 1$.*
- (ii) *If $H^2(G, \mathbb{Z}/p\mathbb{Z}) = H^2(G, \mathbb{Z}/p\mathbb{Z})^+$, then G_Δ is a p -Demuškin group of rank $m + 2$, $0 \leq m \leq n$, with invariant q and dualizing module I .*

Proof: We start with the following remark. Since $\text{Aut}(I) \cong \mathbb{Z}_p^\times$ is abelian, the surjection $G \twoheadrightarrow \Gamma$ factors through G_Δ . With the notation of the proof of proposition 1.1, it follows that $N \cap G$ has infinite index in G and therefore $cd_p(N) = cd_p(N \cap G) \leq 1$, cf. [5] chap.II, §7, ex.3. Using the Hochschild-Serre spectral sequence and the fact that $\text{Hom}(N, \mathbb{Z}/p\mathbb{Z}) = 0$, we see that inf_2 is an isomorphism, and so

$$H^2(G_\Delta) \cong H^2(G)^\Delta.$$

(i) By proposition 1.1, G_Δ is a free pro- p -group. Since the non-degenerate pairing

$$H^1(G) \times H^1(G) \xrightarrow{\cup} H^2(G) \cong \mathbb{Z}/p\mathbb{Z}$$

is Δ -invariant, it follows that

$$\dim_{\mathbb{F}_p} H^1(G)^\pm = n/2 + 1.$$

Therefore

$$\dim_{\mathbb{F}_p} H^1(G_\Delta) = \dim_{\mathbb{F}_p} H^1(G)^\Delta = n/2 + 1.$$

(ii) If $H^2(G) = H^2(G)^+$, then $H^2(G_\Delta) \cong H^2(G)$, and we obtain a non-degenerate pairing

$$H^1(G_\Delta) \times H^1(G_\Delta) \xrightarrow{\cup} H^2(G_\Delta) \cong \mathbb{Z}/p\mathbb{Z}$$

showing that G_Δ is a p -Demuškin group. Finally, since G_Δ is non-trivial and its rank has to be even, it follows that $\dim_{\mathbb{F}_p} H^1(G_\Delta) \geq 2$, and since $\ker(G \twoheadrightarrow G_\Delta)$ acts trivially on I , we have $\#(I^{G_\Delta}) = \#(I^G) = q$ and I is also the dualizing module of G_Δ . $\square$

2 Free Quotients

As before, let G be a p -Demuškin group of rank $n + 2$ with dualizing module I and assume that $2 < q < \infty$. We are interested in quotients of G which are free pro- p -groups. First we calculate the possible ranks of such quotients.

Proposition 2.1 *Let G be a Demuškin group of rank $n + 2$ with finite invariant $q > 2$ and let F be a free quotient of G . Then*

- (i) $H^1(F, \mathbb{Z}/q\mathbb{Z})$ lies in the kernel of the Bockstein homomorphism and
- (ii) $H^1(F, \mathbb{Z}/q\mathbb{Z})$ is a totally isotropic free $\mathbb{Z}/q\mathbb{Z}$ -submodule of $H^1(G, \mathbb{Z}/q\mathbb{Z})$ with respect to the pairing given by the cup-product.

In particular,

$$\text{rank } F \leq \frac{n}{2} + 1.$$

Proof: Since F is free, $H^1(F, \mathbb{Z}/q\mathbb{Z})$ is a free $\mathbb{Z}/q\mathbb{Z}$ -module. The commutative diagram

$$\begin{array}{ccc} H^1(G, \mathbb{Z}/q\mathbb{Z}) \times H^1(G, \mathbb{Z}/q\mathbb{Z}) & \cup & H^2(G, \mathbb{Z}/q\mathbb{Z}) \\ (\text{inf}, \text{inf}) & & \text{inf} \\ H^1(F, \mathbb{Z}/q\mathbb{Z}) \times H^1(F, \mathbb{Z}/q\mathbb{Z}) & \cup & H^2(F, \mathbb{Z}/q\mathbb{Z}) = 0 \end{array}$$

shows that $H^1(F, \mathbb{Z}/q\mathbb{Z})$ is a totally isotropic $\mathbb{Z}/q\mathbb{Z}$ -submodule of $H^1(G, \mathbb{Z}/q\mathbb{Z})$, and so $\dim_{\mathbb{F}_p} H^1(F, \mathbb{Z}/p\mathbb{Z}) = \text{rank}_{\mathbb{Z}/q\mathbb{Z}} H^1(F, \mathbb{Z}/q\mathbb{Z}) \leq n/2 + 1$. From the commutative diagram

$$\begin{array}{ccc} H^1(G, \mathbb{Z}/q\mathbb{Z}) & \xrightarrow{B} & H^2(G, \mathbb{Z}/q\mathbb{Z}) \\ \text{inf} & & \text{inf} \\ H^1(F, \mathbb{Z}/q\mathbb{Z}) & \xrightarrow{B} & H^2(F, \mathbb{Z}/q\mathbb{Z}) = 0 \end{array}$$

follows that $H^1(F, \mathbb{Z}/q\mathbb{Z}) \subseteq \ker(B)$. □

Recall that Γ is the canonical quotient $G/\ker(\chi)$ of G , where $\chi : G \longrightarrow \text{Aut}(I)$ is the character given by the action of G on I , i.e.

$$\chi_0 : \Gamma \longrightarrow \text{Aut}(I).$$

Lemma 2.2 *The submodules $H^1(\Gamma, \mathbb{Z}/q\mathbb{Z})$ and $\ker B$ of $H^1(G, \mathbb{Z}/q\mathbb{Z})$ are orthogonal to each other, more precisely*

$$H^1(\Gamma, \mathbb{Z}/q\mathbb{Z}) = (\ker B)^\perp.$$

Proof: Consider the commutative diagram of non-degenerate pairings

$$\begin{array}{ccc} H^1(G, {}_qI) \times H^1(G, \mathbb{Z}/q\mathbb{Z}) & \cup & H^2(G, {}_qI) \quad \mathbb{Z}/q\mathbb{Z} \\ \delta & & B \\ H^0(G, {}_qI) \times H^2(G, \mathbb{Z}/q\mathbb{Z}) & \cup & H^2(G, {}_qI) \quad \mathbb{Z}/q\mathbb{Z} \end{array}$$

which is induced by the exact sequences

$$0 \longrightarrow \mathbb{Z}/q\mathbb{Z} \xrightarrow{q} \mathbb{Z}/q^2\mathbb{Z} \longrightarrow \mathbb{Z}/q\mathbb{Z} \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow {}_qI \xrightarrow{q} {}_{q^2}I \longrightarrow {}_qI \longrightarrow 0.$$

Let ζ_{q^2} be a generator of ${}_{q^2}I$, and so $\zeta_q = (\zeta_{q^2})^q$ is a generator of ${}_qI$. By definition of the homomorphism δ , i.e.

$$\delta : H^0(G, {}_qI) \longrightarrow H^1(G, {}_qI), \quad (\zeta_q)^i \mapsto \left\{ G \rightarrow {}_qI, \quad g \mapsto ((\zeta_{q^2})^i)^{g-1} \right\},$$

we see that the image of δ is $H^1(\Gamma, {}_qI)$. Since the pairings above are non-degenerated, it follows that $H^1(\Gamma, \mathbb{Z}/q\mathbb{Z}) = H^1(\Gamma, {}_qI)$ is orthogonal to $\ker B$, and since $\text{rank}_{\mathbb{Z}/q\mathbb{Z}}(\ker B)^\perp = 1$, we prove the lemma. $\square$

Proposition 2.3 *Let G be a p -Demuškin group of rank $n+2$ with finite invariant $q > 2$ and let F be a free factor of G of rank $n/2+1$. Then the canonical surjection $G \twoheadrightarrow \Gamma$ factors through F , i.e. there is a commutative diagram*

$$\begin{array}{ccc} G & & \Gamma \\ & \searrow & \uparrow \\ & F & \end{array}$$

Proof: Suppose the contrary. Then there exists an open subgroup G' of G which has a surjection

$$(G')^{ab} \twoheadrightarrow (F')^{ab} \times \Gamma',$$

where F' is the image of G' in F under the projection $G \twoheadrightarrow F$ and Γ' is the image of G' under the projection $G \twoheadrightarrow \Gamma$. Let $q' = \#(I^{G'}) = \#(I^{\Gamma'})$. Since F' is free, it follows that $H^1(F', \mathbb{Z}/q'\mathbb{Z})$ is a totally isotropic submodule of $H^1(G', \mathbb{Z}/q'\mathbb{Z})$ and contained in $\ker B'$ by proposition 2.1. From lemma 2.2 we know that $H^1(\Gamma', \mathbb{Z}/q'\mathbb{Z})$ is orthogonal to $\ker B'$, and so also to $H^1(F', \mathbb{Z}/q'\mathbb{Z})$. Thus $H^1(F', \mathbb{Z}/q'\mathbb{Z}) \oplus H^1(\Gamma', \mathbb{Z}/q'\mathbb{Z})$ is totally isotropic. But $H^1(F', \mathbb{Z}/q'\mathbb{Z})$ is a maximal totally isotropic $\mathbb{Z}/q'\mathbb{Z}$ -submodule of $H^1(G', \mathbb{Z}/q'\mathbb{Z})$ of rank $d \cdot n/2 + 1$, where $d = (G : G')$. This contradiction proves the proposition. $\square$

For the existence of free quotients of Demuškin groups we have the following

Theorem 2.4 *Let G be a p -Demuškin group of rank $n + 2$ with finite invariant $q > 2$ and let Δ be a finite abelian group of exponent $p - 1$ acting on G . Let V be a $\mathbb{Z}/q\mathbb{Z}$ -submodule of $H^1(G, \mathbb{Z}/q\mathbb{Z})$ such that*

- (i) V is $\mathbb{Z}/q\mathbb{Z}$ -free and Δ -invariant,
- (ii) V is totally isotropic with respect to the pairing given by the cup-product,
- (iii) V lies in the kernel of the Bockstein map $B : H^1(G, \mathbb{Z}/q\mathbb{Z}) \rightarrow H^2(G, \mathbb{Z}/q\mathbb{Z})$.

Then there exists a Δ -invariant surjection

$$G \twoheadrightarrow F$$

onto a free quotient F of G such that $H^1(F, \mathbb{Z}/q\mathbb{Z}) = V$.

Proof: Let

$$1 \longrightarrow R \longrightarrow F_{n+2} \longrightarrow G \longrightarrow 1$$

be a minimal presentation of G , where F_{n+2} is a free pro- p -group of rank $n + 2$. Using lemma 1.2, we extend the action of Δ to F_{n+2} . Let $\gamma, x_0, \dots, x_n$ be a basis of F_{n+2} such that

- (i) each element of the basis of F_{n+2} generates a Δ -invariant subgroup isomorphic to $\mathbb{Z}_p$ on which Δ acts by some character $\psi : \Delta \rightarrow \mu_{p-1}$,
- (ii) R , as a normal subgroup of F_{n+2} , is generated by the element

$$w = (x_0)^q [x_0, \gamma] [x_1, x_2] [x_3, x_4] \cdots [x_{n-1}, x_n] \cdot f$$

where $f \in (F_{n+2})^3$,

- (iii) $V^* = \text{Hom}(V, \mathbb{Z}/q\mathbb{Z})$ has a basis $\{v_i \bmod (F_{n+2})^2, 1 \leq i \leq r = \text{rank}_{\mathbb{Z}/q\mathbb{Z}} V\}$ such that

$$\{v_1, \dots, v_r\} \quad \text{is a subset of} \quad \{\gamma, x_1, \dots, x_n\}$$

and, if $v_i = x_{j(i)}$, then $x_{j(i)+1} \notin \{v_1, \dots, v_r\}$ or $x_{j(i)-1} \notin \{v_1, \dots, v_r\}$ according to whether $j(i)$ is odd or even.

Such a basis exists: by [1] prop.(2.3), we find a basis of F_{n+2} with the property (i) (since Δ is abelian of exponent $p - 1$, the $\mathbb{Z}/q\mathbb{Z}[\Delta]$ -module $F_{n+2}/(F_{n+2})^2$ decomposes into a sum of $\mathbb{Z}/q\mathbb{Z}[\Delta]$ -modules which are free $\mathbb{Z}/q\mathbb{Z}$ -modules of rank equal to 1). The Δ -invariance of the cup-product and the Bockstein homomorphism implies that we find a basis satisfying (i) and (ii), cf. [8] lemma 3 in the case where $\Delta \cong \mathbb{Z}/2\mathbb{Z}$. Using the assumptions on V , we can also satisfy (iii).

Let N be the normal subgroup of F_{n+2} generated by the set

$$\{\gamma, x_k, 0 \leq k \leq n\} \setminus \{v_1, \dots, v_r\},$$

then $F := F_{n+2}/N$ is a free pro- p -group of rank r , N is Δ -invariant and we have $R \subseteq N(F_{n+2})^3$ by the properties (ii) and (iii) of the basis $\gamma, x_0, \dots, x_n$. Thus the Δ -invariant surjection

$$F_{n+2} \twoheadrightarrow F/F^3 = F_{n+2}/N(F_{n+2})^3$$

factors through a Δ -invariant surjection

$$G \twoheadrightarrow F/F^3.$$

Applying proposition 1.3, we get a Δ -invariant surjection from G onto a free pro- p -group F which induces a surjection $G \twoheadrightarrow F/F^2 \cong F_{n+2}/N(F_{n+2})^2$.

By construction, we have $F/F^2 \cong V^*$, and so $H^1(F, \mathbb{Z}/q\mathbb{Z}) = V$. This finishes the proof of the theorem. $\square$

Now we consider free quotients of a Demuškin group G which are invariant under a given Δ -action of G , where Δ is a group of order 2.

Corollary 2.5 *Let p be an odd prime number and let G be a p -Demuškin group of rank $n + 2$, $n \geq 0$, with finite invariant q . Let $\Delta \cong \mathbb{Z}/2\mathbb{Z}$ acting on G such that $H^2(G, \mathbb{Z}/q\mathbb{Z}) = H^2(G, \mathbb{Z}/q\mathbb{Z})^-$. Let*

$$u^+, u^- \geq 0 \quad \text{be integers such that} \quad u^+ + u^- = n/2.$$

Then there exists a Δ -invariant surjection

$$\varphi : G \twoheadrightarrow F$$

such that

- (i) F is a free pro- p -group of rank $n/2 + 1$,
- (ii) $\text{rank}_{\mathbb{Z}_p}(F^{ab})^+ = u^+ + 1$ and $\text{rank}_{\mathbb{Z}_p}(F^{ab})^- = u^-$.

Proof: Since $H^2(G, \mathbb{Z}/q\mathbb{Z}) = H^2(G, \mathbb{Z}/q\mathbb{Z})^-$, the submodules $H^1(G, \mathbb{Z}/q\mathbb{Z})^\pm$ are maximal totally isotropic with respect to the cup-product pairing, and so

$$\text{rank}_{\mathbb{Z}/q\mathbb{Z}} H^1(G, \mathbb{Z}/q\mathbb{Z})^\pm = n/2 + 1.$$

Let

$$V = V^+ \oplus V^-$$

where V^+ is a free $\mathbb{Z}/q\mathbb{Z}$ -submodule of $H^1(G, \mathbb{Z}/q\mathbb{Z})^+$ of rank $1 + u^+$ containing $H^1(\Gamma, \mathbb{Z}/q\mathbb{Z})$, and V^- is a free $\mathbb{Z}/q\mathbb{Z}$ -submodule of $\ker B^-$ of rank u^- being orthogonal to V^+ , i.e. V is maximal totally isotropic and contained in $\ker B$.

It is easy to see that V^- exists (see also the remark below): by lemma 2.2 $H^1(\Gamma, \mathbb{Z}/q\mathbb{Z}) \subseteq (\ker B^-)^\perp$, and since

$$\text{rank}_{\mathbb{Z}/q\mathbb{Z}} H^1(G, \mathbb{Z}/q\mathbb{Z})^+ - \text{rank}_{\mathbb{Z}/q\mathbb{Z}} V^+ = n/2 + 1 - (1 + u^+) = u^-,$$

there exists a free submodule of $\ker B^-$ of rank u^- which is orthogonal to V^+ . Now theorem 2.4 gives us a free Δ -invariant quotient F of G of rank $n/2 + 1$ such that

$$H^1(F, \mathbb{Z}/q\mathbb{Z}) = V \cong (\mathbb{Z}/q\mathbb{Z}[\Delta]^+)^{u^++1} \oplus (\mathbb{Z}/q\mathbb{Z}[\Delta]^-)^{u^-}.$$

Since F^{ab} is free $\mathbb{Z}_p$ -module, we obtain assertion (ii). $\square$

Remark: Explicitly, we get a submodule V with the properties as above in the following way: let

$$1 \longrightarrow R \longrightarrow F_{n+2} \longrightarrow G \longrightarrow 1$$

be a minimal presentation of G , where F_{n+2} is a free pro- p -group of rank $n+2$ with the extended action of Δ . Let $\gamma, x_0, \dots, x_n$ be a basis of F_{n+2} such that R is generated by the element

$$w = (x_0)^q [x_0, \gamma] [x_1, x_2] [x_3, x_4] \cdots [x_{n-1}, x_n] \cdot f,$$

$f \in (F_{n+2})^3$, and

$$\begin{aligned} \gamma^\sigma &= \gamma \cdot a, & x_i^\sigma &= x_i \cdot a_i & \text{for } i = 2, 4, \dots, n, \\ x_0^\sigma &= x_0^{-1} \cdot b, & x_i^\sigma &= x_i^{-1} \cdot b_i & \text{for } i = 1, 3, 5, \dots, n-1. \end{aligned}$$

with $a, b, a_i, b_i \in (F_{n+2})^2$. Such a basis exists by the Δ -invariance of the cup-product and the Bockstein homomorphism, cf. [8] lemma 3. If we put

$$\begin{aligned} \gamma' &:= \gamma \cdot a^{\frac{1}{2}}, & x'_i &:= x_i \cdot a_i^{\frac{1}{2}} & \text{for } i = 2, 4, \dots, n, \\ x'_0 &:= b^{-\frac{1}{2}} \cdot x_0, & x'_i &:= b_i^{-\frac{1}{2}} \cdot x_i & \text{for } i = 1, 3, 5, \dots, n-1, \end{aligned}$$

then

$$\begin{aligned} (\gamma')^\sigma &= \gamma', & (x'_i)^\sigma &= x'_i & \text{for } i \geq 2 \text{ even}, \\ (x'_0)^\sigma &= (x'_0)^{-1}, & (x'_i)^\sigma &= (x'_i)^{-1} & \text{for } i \geq 1 \text{ odd}, \end{aligned}$$

and

$$w = (x'_0)^q [x'_0, \gamma'] [x'_1, x'_2] [x'_3, x'_4] \cdots [x'_{n-1}, x'_n] \cdot f'$$

where $f' \in (F_{n+2})^3$. Let $u = 2u^+ - 1$. If we denote $x \bmod F^2$ by $\bar{x}$, then the dual of

$$\begin{aligned} V^* &:= \mathbb{Z}/q\mathbb{Z} \cdot \bar{\gamma} \oplus \bigoplus_{i=1,3,\dots,u} \mathbb{Z}/q\mathbb{Z} \cdot \bar{x}_{i+1} \oplus \bigoplus_{i=u+3,\dots,n} \mathbb{Z}/q\mathbb{Z} \cdot \bar{x}_{i-1} \\ &\cong (\mathbb{Z}/q\mathbb{Z}[\Delta]^+)^{u^++1} \oplus (\mathbb{Z}/q\mathbb{Z}[\Delta]^-)^{u^-} \end{aligned}$$

gives an example for a submodule with the properties (i)-(iii) in the proof of corollary 2.5. The free quotient of G is obtained in the following way: if

$$N' = (\underbrace{x'_0, x'_1, x'_3, \dots, x'_u}_{u^+ \text{-times}}, \underbrace{x'_{u+3}, \dots, x'_n}_{u^- \text{-times}}) \trianglelefteq F_{n+2},$$

then $F = F_{n+2}/N'$ is a free pro- p -group of rank $n/2 + 1$, N' is Δ -invariant, $R \subseteq N(F_{n+2})^3$ and $V^* = F/F^2$. Using proposition 1.3 we get the desired quotient of G .

With the notation and assumptions of corollary 2.5, we make for a Δ -invariant free quotient F of G of rank $n/2 + 1$ the following

Definition 2.6 *We call the tuple (u^+, u^-) the **signature** of F , if*

$$F/F^2 \cong (\mathbb{Z}/q\mathbb{Z}[\Delta]^+)^{u^++1} \oplus (\mathbb{Z}/q\mathbb{Z}[\Delta]^-)^{u^-}.$$

It is easy to see that the signature of a maximal free quotient F of G does not determine F , but if the signature is equal to $(n/2, 0)$, then we have the following proposition.

Proposition 2.7 *Let p be an odd prime number and let Δ be of order 2. Let G be a p -Demuškin group of rank $n + 2$ with finite invariant q on which Δ acts such that $H^2(G, \mathbb{Z}/p\mathbb{Z})^\Delta = 0$. Let F be a free Δ -invariant quotient of G of rank $n/2 + 1$, i.e. the canonical surjection*

$$G \twoheadrightarrow F$$

is Δ -invariant. If the induced action of Δ on F/F^2 is trivial, i.e. F has signature $(n/2, 0)$, then F is equal to the maximal quotient G_Δ of G with trivial Δ -action. In particular, a free quotient of G with the properties above is unique.

Proof: As in the remark after the proof of corollary 2.5, we find generators of F on which Δ acts trivially, and so F has a trivial Δ -action. Thus we have a surjection

$$\varphi : G_\Delta \twoheadrightarrow F.$$

Since G_Δ is free of rank $n/2 + 1 = \dim_{\mathbb{F}_p} H^1(F, \mathbb{Z}/p\mathbb{Z})$ by proposition 1.4(i), it follows that φ is an isomorphism. Thus F is the maximal quotient of G with trivial Δ -action. $\square$

References

- [1] Boston, N. *Explicit deformation of Galois representations*. Invent. math. **103** (1991) 181-196
- [2] Jannsen, U., Wingberg, K. *Einbettungsprobleme und Galoisstruktur lokaler Körper*. J. reine u. angew. Math. **319** (1980) 196-212
- [3] Labute, J.,P. *Classification of Demuškin groups*. Can. J. Math. **19** (1967) 106-132
- [4] Lubotzky, A. *Combinatorial group theory for pro-p-groups*. J. of Pure and Appl. Algebra **25** (1982) 311-325
- [5] Neukirch, J., Schmidt, A., Wingberg, K. *Cohomology of Number Fields*. Springer 1999
- [6] Nguyen-Quand-Do, T. *Sur la structure galoisienne des corps locaux et la théorie d'Iwasawa*. Compos. Math. **46** (1982) 85-119
- [7] Sonn, J. *Epimorphisms of Demushkin Groups*. Israel J. Math. **17** (1974) 176-190
- [8] Wingberg, K. *On Demuškin groups with involution*. Ann. Sci. Éc. Norm. Sup. 4^e série **22** (1989) 555-567

Mathematisches Institut
der Universität Heidelberg
Im Neuenheimer Feld 288
69120 Heidelberg
Germany

e-mail: wingberg@mathi.uni-heidelberg.de